

Comments on Bosonization of the extended Thirring Model with $SU(2)$ Symmetry

Jiro SAKAMOTO^{*)} and Keiji MANO

Department of Material Science, Shimane University, Matsue 690-8504, Japan

Bosonization of the extended Thirring model with $SU(2)$ symmetry in the Minkowski path integral method is discussed. We argue that it is not an easy task to bosonize such a model if we derive correctly the fermion determinant which is induced with the decoupling transformation because it seems that there arise ghost fields. This is contrary to what is shown in a textbook and some people believe.

§1. Introduction

In a previous paper ¹⁾ we derived the direct correspondence between the fermion currents and the boson fields for the extended Thirring model with $SU(2)$ symmetry in the Minkowski path integral method, and we showed that the corresponding boson model contains ghost fields which interact with the other boson fields in a complex manner and that it is hard to eliminate them, while in the original Thirring model such ghost field also arises but it is decoupled from other fields and disappears in the final Lagrangian.

Since Coleman carried out the bosonization of the massive Thirring model and showed that it is equivalent to the sine-Gordon model, ²⁾ many people have attempted to apply this technique to fermion models with non-Abelian symmetries, ³⁾ and some people believe that the bosonization of the Thirring model with $SU(N)$ symmetry can be easily carried out. ⁴⁾ Their consequences, however, do not agree with ours mentioned above. In the present paper we argue that in their formulation the fermion determinant, which is induced by the decoupling transformation of the fermion fields, is not correctly derived.

In the next section, after reviewing the formulation of Ref. 4) briefly we derive the correct expression of the fermion determinant for the decoupling transformation following Fujikawa's method. ^{3), 5), 6)} We use the same notation as in the previous paper. ¹⁾: $g_{\mu\nu} = (-1, +1)$, $\epsilon^{01} = -\epsilon_{01} = 1$, $\gamma^0 = i\sigma_x$, $\gamma^1 = \sigma_y$ and $\gamma^5 = \sigma_z$. The 2×2 matrices of the $SU(2)$ generators are given by $\tau_1 = \frac{1}{2}\sigma_x$, $\tau_2 = \frac{1}{2}\sigma_y$ and $\tau_3 = \frac{1}{2}\sigma_z$. We use the first part of the Roman alphabet as $SU(2)$ suffixes 1, 2, 3 and the middle part i, j as only 1 and 2.

^{*)} E-mail: jsakamot@riko.shimane-u.ac.jp

§2. Derivation of the fermion determinant

Our initial Lagrangian for the extended Thirring model with $SU(2)$ symmetry is given by

$$\mathcal{L} = \bar{\psi}\not{\partial}\psi + \frac{g}{2} \sum_{a=1}^3 j_\mu^a j_\mu^a + \frac{g'}{2} j_\mu j_\mu, \quad (2.1)$$

with the $SU(2)$ current

$$j_\mu^a = i\bar{\psi}\tau_a\gamma_\mu\psi \quad (2.2)$$

and the $U(1)$ current

$$j_\mu = i\bar{\psi}\gamma_\mu\psi. \quad (2.3)$$

The last term in Eq.(2.1) is necessary for renormalizability of the model. The generating functional is given by

$$Z = \int d\bar{\psi}d\psi \exp i \int d^2x \mathcal{L}. \quad (2.4)$$

Now we review the formulation of Ref. 4) briefly. We divide Eq.(2.1) into two parts as follows,

$$\mathcal{L}_0 = \bar{\psi}\not{\partial}\psi + \frac{g}{2} j_\mu^3 j_\mu^3 + \frac{g'}{2} j_\mu j_\mu, \quad (2.5)$$

$$\mathcal{L}_I = \frac{g}{2} \sum_{i=1}^2 j_\mu^i j_\mu^i. \quad (2.6)$$

$\mathcal{L}_0$ of the above expressions consists of the diagonal parts of the $SU(2)$ multiplets and is treated as the non-perturbative part of the Lagrangian, and $\mathcal{L}_I$ is treated as the perturbative part. Eq.(2.5) is equivalently rewritten as

$$\mathcal{L}_0 = \bar{\psi}\not{\partial}\psi + j_\mu^3 A_\mu^3 - \frac{1}{2g} A_\mu^3 A_\mu^3 + j_\mu a^\mu - \frac{1}{2g'} a_\mu a^\mu, \quad (2.7)$$

with auxiliary vector fields A_μ^3 and a_μ . In two-dimensional spacetime, these vector fields are expressed as

$$A_\mu^3 = \partial_\mu \Lambda_3 + \epsilon_{\mu\nu} \partial^\nu \Phi_3, \quad (2.8)$$

$$a_\mu = \partial_\mu \lambda + \epsilon_{\mu\nu} \partial^\nu \phi. \quad (2.9)$$

We substitute the above expressions into Eq.(2.7) to obtain

$$\begin{aligned} \mathcal{L}_0 = & \bar{\psi}\{\not{\partial} + i\tau_3(\not{\partial}\Lambda_3 + \gamma_5\not{\partial}\Phi_3) + i(\not{\partial}\lambda + \gamma_5\not{\partial}\phi)\}\psi \\ & - \frac{1}{2g}\{\partial^\mu \Lambda_3 \partial_\mu \Lambda_3 - \partial^\mu \Phi_3 \partial_\mu \Phi_3\} - \frac{1}{2g'}\{\partial^\mu \lambda \partial_\mu \lambda - \partial^\mu \phi \partial_\mu \phi\}, \end{aligned} \quad (2.10)$$

where we use the identity $\gamma_5\gamma_\mu = -\epsilon_{\mu\nu}\gamma^\nu$. The generating functional for $\mathcal{L}_0$ is given by

$$Z_0 = \int d\bar{\psi}d\psi d\Lambda_3 d\Phi_3 d\lambda d\phi \exp i \int d^2x \mathcal{L}_0. \quad (2.11)$$

The coupling terms between the fermion and boson fields in (2·10) are eliminated by the following decoupling transformation,

$$\psi \rightarrow \psi' = U(\Lambda_3, \Phi_3, \lambda, \phi)\psi, \quad (2\cdot12a)$$

$$U(\Lambda_3, \Phi_3, \lambda, \phi) = \exp i\{\tau_3(\Lambda_3 - \gamma_5\Phi_3) + \lambda - \gamma_5\phi\}, \quad (2\cdot12b)$$

and we have

$$\mathcal{L}_0 = \overline{\psi'}\not{\partial}\psi' - \frac{1}{2g}\{\partial^\mu\Lambda_3\partial_\mu\Lambda_3 - \partial^\mu\Phi_3\partial_\mu\Phi_3\} - \frac{1}{2g'}\{\partial^\mu\lambda\partial_\mu\lambda - \partial^\mu\phi\partial_\mu\phi\}. \quad (2\cdot13)$$

The kinetic terms of Φ_3 and ϕ in the above expression have wrong signs and they seem to be ghost fields. As is seen in the following, however, ϕ is decoupled from the other fields and Φ_3 is absorbed to the other fields and they finally disappear.

The fermion Jacobian which is induced with the transformation eq.(2·12) is derived following Fujikawa's method as^{1), 3), 5)}

$$J = \exp i \int d^2x \frac{1}{4\pi} \{(\partial\Phi_3)^2 + 2(\partial\phi)^2\}, \quad (2\cdot14)$$

where we use the eigenfunction $\psi_n(x; t)$ defined by

$$i\not{D}(t)\psi_n(x; t) \equiv i(\not{\partial} + i\not{\phi}(t))\psi_n(x; t) = \lambda_n\psi_n(x; t) \quad (2\cdot15)$$

as the basis functions for the Jacobian. Here we put

$$\begin{aligned} \not{\phi}(t) &= U(-\Lambda_3, \Phi_3, -\lambda, \phi; -t)\not{\phi}U(\Lambda_3, \Phi_3, \lambda, \phi; -t) \\ &\quad - iU(-\Lambda_3, \Phi_3, -\lambda, \phi; -t)\not{\partial}U(\Lambda_3, \Phi_3, \lambda, \phi; -t), \end{aligned} \quad (2\cdot16)$$

with

$$\not{\phi} = A_3^\mu\gamma_\mu\tau_3 + a^\mu\gamma_\mu = (\not{\partial}\Lambda_3 + \gamma_5\not{\partial}\Phi_3)\tau_3 + \not{\partial}\lambda + \gamma_5\not{\partial}\phi, \quad (2\cdot17)$$

$$U(\Lambda_3, \Phi_3, \lambda, \phi; t) = \exp it\{\tau_3(\Lambda_3 - \gamma_5\Phi_3) + \lambda - \gamma_5\phi\}. \quad (2\cdot18)$$

Eq.(2·15) corresponds to the Euler equation for the fermion field obtained from $\mathcal{L}_0$ of (2·7),

$$i(\not{\partial} + i\not{\phi})\psi = 0. \quad (2\cdot19)$$

By the decoupling transformation (2·12), currents j_μ^i ($i = 1, 2$) in $\mathcal{L}_I$ of (2·6) are transformed as

$$j_\mu^i = i\overline{\psi}\tau_i\gamma_\mu\psi = i\overline{\psi'}\tau_i\gamma_\mu\exp\{-2i\tau_3(\Lambda_3 - \gamma_5\Phi_3)\}\psi', \quad (2\cdot20)$$

and $\mathcal{L}_I$ is rewritten as

$$\mathcal{L}_I = -\frac{g}{2}\overline{\psi'}_1\gamma_\mu\exp\{-i\gamma_5\Phi_3\}\psi'_2\overline{\psi'}_2\gamma^\mu\exp\{i\gamma_5\Phi_3\}\psi'_1, \quad (2\cdot21)$$

where we put ψ' of $SU(2)$ -doublet as

$$\psi' = \begin{pmatrix} \psi'_1 \\ \psi'_2 \end{pmatrix}. \quad (2\cdot22)$$

Eq.(2.21) is rewritten as

$$\begin{aligned}\mathcal{L}_I &= g \left\{ \overline{\psi}'_{1+} \psi'_{2+} \overline{\psi}'_{2-} \psi'_{1-} \exp\{-2i\Phi_3\} + \overline{\psi}'_{1-} \psi'_{2-} \overline{\psi}'_{2+} \psi'_{1+} \exp\{2i\Phi_3\} \right\} \\ &= -g \left\{ \overline{\psi}'_{1+} \psi'_{1-} \overline{\psi}'_{2-} \psi'_{2+} \exp\{-2i\Phi_3\} + \overline{\psi}'_{1-} \psi'_{1+} \overline{\psi}'_{2+} \psi'_{2-} \exp\{2i\Phi_3\} \right\}\end{aligned}\quad (2.23)$$

with putting spin-doublet as

$$\psi'_i = \begin{pmatrix} \psi'_{i+} \\ \psi'_{i-} \end{pmatrix}. \quad (2.24)$$

We note that all the boson fields except Φ_3 are decoupled from the fermion fields and can be eliminated by the path integration. In a perturbative expansion over $\mathcal{L}_I$ we see that we can replace the fermion fields with their boson counterparts θ_i and obtain the effective Lagrangian

$$\begin{aligned}\mathcal{L} &= -\frac{1}{2g} \left\{ (\partial\Lambda_3)^2 - (\partial\Phi_3)^2 \right\} - \frac{1}{2g'} \left\{ (\partial\lambda)^2 - (\partial\phi)^2 \right\} + \frac{1}{4\pi} (\partial\Phi_3)^2 + \frac{1}{2\pi} (\partial\phi)^2 \\ &\quad - \frac{1}{2} (\partial\theta_1)^2 - \frac{1}{2} (\partial\theta_2)^2 + g\mu^2 \cos \left\{ \sqrt{4\pi}(\theta_1 + \theta_2) + 2\Phi_3 \right\},\end{aligned}\quad (2.25)$$

where μ is an infrared cutoff parameter with mass dimension.^{4), 6)} We put

$$\Theta = \sqrt{4\pi}(\theta_1 + \theta_2) + 2\Phi_3, \quad (2.26)$$

to change $\Phi_3 \rightarrow \Theta$ and integrate over θ_i, λ, ϕ and Λ_3 to obtain

$$\mathcal{L} = -\frac{1}{16\pi} \left(1 + \frac{g}{2\pi} \right) (\partial\Theta)^2 + g\mu^2 \cos \Theta. \quad (2.27)$$

As mentioned before the ghost field ϕ is decoupled from other fields and Φ_3 is absorbed into Θ , and there is no ghost field in the above expression.

The simple expression (2.27), however, obviously disagrees with our previous consequence in Ref.1). The decomposition of the Lagrangian into Eqs.(2.5) and (2.6) violates $SU(2)$ symmetry and the fermion Jacobian (2.14) is not correctly derived. To see this we introduce auxiliary vector fields A_μ^a and a_μ to rewrite (2.1) as

$$\begin{aligned}\mathcal{L} &= \overline{\psi} \not{\partial} \psi + j_\mu^a A_\mu^a - \frac{1}{2g} A_\mu^a A_\mu^a + j_\mu a^\mu - \frac{1}{2g'} a^\mu a_\mu \\ &= \overline{\psi} \{ \not{\partial} + i\tau_a (\not{\partial} \Lambda^a + \gamma^\mu \epsilon_{\mu\nu} \partial^\nu \Phi^a) + i(\not{\partial} \lambda + \gamma^\mu \epsilon_{\mu\nu} \partial^\nu \phi) \} \psi \\ &\quad - \frac{1}{2g} \{ \partial^\mu \Lambda_a \partial_\mu \Lambda_a - \partial^\mu \Phi_a \partial_\mu \Phi_a \} - \frac{1}{2g'} \{ \partial^\mu \lambda \partial_\mu \lambda - \partial^\mu \phi \partial_\mu \phi \},\end{aligned}\quad (2.28)$$

where we put

$$A_\mu^a = \partial_\mu \Lambda_a + \epsilon_{\mu\nu} \partial^\nu \Phi_a, \quad (2.29a)$$

$$a_\mu = \partial_\mu \lambda + \epsilon_{\mu\nu} \partial^\nu \phi. \quad (2.29b)$$

By the decoupling transformation (2.12), Lagrangian (2.28) is transformed into

$$\begin{aligned}\mathcal{L} &= \overline{\psi}' \{ \not{\partial} + i \sum_{i=1,2} A'_i \tau_i \} \psi' \\ &\quad - \frac{1}{2g} \{ \partial^\mu \Lambda_a \partial_\mu \Lambda_a - \partial^\mu \Phi_a \partial_\mu \Phi_a \} - \frac{1}{2g'} \{ \partial^\mu \lambda \partial_\mu \lambda - \partial^\mu \phi \partial_\mu \phi \},\end{aligned}\quad (2.30)$$

where we put

$$\begin{aligned}\sum_{i=1,2} A'_i \tau_i &\equiv \sum_{i=1,2} e^{i\tau_3(\Lambda_3 + \gamma_5 \Phi_3)} A_i \tau_i e^{-i\tau_3(\Lambda_3 - \gamma_5 \Phi_3)} \\ &= \sum_{i=1,2} e^{2i\tau_3(\Lambda_3 + \gamma_5 \Phi_3)} A_i \tau_i.\end{aligned}\quad (2.31)$$

To derive the Jacobian for the decoupling transformation (2.12) we use the eigenfunction $\psi_n(x; t)$ which satisfies

$$i\mathcal{D}'(t)\psi_n(x; t) \equiv i(\not{\partial} + i\mathbf{A}(t))\psi_n(x; t) = \lambda_n \psi_n(x; t), \quad (2.32)$$

instead of (2.15), where we define $\mathbf{A}(t)$ as

$$\begin{aligned}\mathbf{A}(t) &= U(-\Lambda_3, \Phi_3, -\lambda, \phi; -t) \mathbf{A} U(\Lambda_3, \Phi_3, \lambda, \phi; -t) \\ &\quad - iU(-\Lambda_3, \Phi_3, -\lambda, \phi; -t) \not{\partial} U(\Lambda_3, \Phi_3, \lambda, \phi; -t),\end{aligned}\quad (2.33)$$

with

$$\mathbf{A} = \sum_{a=1}^3 A_a \tau_a + \not{\phi}. \quad (2.34)$$

Eq.(2.33) is rewritten as

$$\mathbf{A}(t) = (1-t)(A_3 \tau_3 + \not{\phi}) + \exp\{2it\tau_3(\Lambda_3 + \gamma_5 \Phi_3)\} \sum_{i=1,2} A_i \tau_i. \quad (2.35)$$

We see $\mathbf{A}(1) = \sum_{i=1,2} A'_i \tau_i$. The equation (2.32) corresponds to the Euler equation for the fermion field obtained from the Lagrangian (2.28);

$$i(\not{\partial} + i\mathbf{A})\psi = 0. \quad (2.36)$$

The fermion measure is transformed under (2.12) as

$$d\psi = |\det U(\Lambda_3, \Phi_3, \lambda, \phi)| d\psi', \quad (2.37)$$

$$d\bar{\psi} = |\det U(-\Lambda_3, \Phi_3, -\lambda, \phi)| d\bar{\psi}'. \quad (2.38)$$

The determinant of the above expression (2.37) is derived as ¹⁾

$$|\det U(\Lambda_3, \Phi_3, \lambda, \phi)| = \exp \frac{i}{4\pi} \int d^2x W, \quad (2.39)$$

with

$$\begin{aligned}W &= \int_0^1 dt \text{Tr} [\gamma_5(\phi + \tau_3 \Phi_3) \{ \not{\partial} \mathbf{A}(t) + i\mathbf{A}(t) \mathbf{A}(t) \}] \\ &= \int_0^1 dt \text{Tr} [\gamma_5(\phi + \tau_3 \Phi_3) \{ -(1-t) \epsilon_{\mu\nu} \gamma_5 (\partial^\mu A_3^\nu \tau_3 + \partial^\mu a^\nu) \\ &\quad + \tau_3 \gamma_5 \epsilon_{ij} \epsilon_{\mu\nu} A_i^\mu A_j^\nu \cos(2t\Phi_3) - \tau_3 \gamma_5 A_i^\nu A_j^\nu \delta_{ij} g_{\mu\nu} \sin(2t\Phi_3) \}] \\ &= - \left[\frac{1}{2} \epsilon_{\mu\nu} \partial^\mu A_3^\nu \Phi_3 + \epsilon_{\mu\nu} \partial^\mu a^\nu \phi \right]\end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2}\epsilon_{ij}\epsilon_{\mu\nu}A_i^\mu A_j^\nu \sin(2\Phi_3) + \frac{1}{2}g_{\mu\nu}A_i^\mu A_i^\nu \{1 - \cos(2\Phi_3)\} \Big] \\
& = \left[\frac{1}{2}(\partial\Phi_3)^2 + (\partial\phi)^2 + \epsilon_{ij}\partial^\mu A_i \partial_\mu \Phi_j \sin(2\Phi_3) \right. \\
& \quad \left. - \frac{1}{2}(\partial^\mu A_i \partial_\mu A_i - \partial^\mu \Phi_i \partial_\mu \Phi_i) \{1 - \cos(2\Phi_3)\} \right]. \tag{2.40}
\end{aligned}$$

The determinant (2.38) is derived samely. Then the effective Lagrangian including these determinants is given by

$$\begin{aligned}
\mathcal{L} = & \bar{\psi}'(\not{\partial} + iA'_i\tau_i)\psi' - \frac{1}{2g} \left\{ (\partial A_a)^2 - (\partial \Phi_a)^2 \right\} - \frac{1}{2g'} \left\{ (\partial \lambda)^2 - (\partial \phi)^2 \right\} \\
& + \frac{1}{4\pi} \left[(\partial\Phi_3)^2 + 2(\partial\phi)^2 \right. \\
& \left. + 2\epsilon_{ij}\partial^\mu A_i \partial_\mu \Phi_j \sin(2\Phi_3) - (\partial^\mu A_i \partial_\mu A_i - \partial^\mu \Phi_i \partial_\mu \Phi_i) \{1 - \cos(2\Phi_3)\} \right]. \tag{2.41}
\end{aligned}$$

We can obtain the same effective Lagrangian as (2.25) by the path integration over A_i and Φ_i if the last line in the above equation does not exist.

§3. Concluding remark

We discuss bosonization of the extended Thirring model with $SU(2)$ symmetry. As in the textbook of Ref.4), we divide the current-current interaction $g \sum_{a=1}^3 j_a^\mu j_\mu^a + g' j^\mu j_\mu$ into $gj_3^\mu j_\mu^3 + g' j^\mu j_\mu$ and $g \sum_{a=1}^2 j_a^\mu j_\mu^a$. We convert the former interaction terms into $j_\mu^3 A_3^\mu + j_\mu a^\mu$ with introducing the auxiliary boson fields A_μ^3 and a_μ and eliminate them from the action by the decoupling transformation. Calculating the Jacobian induced by the decoupling transformation and perturbative expansion with respect to the latter interaction terms, we obtain the corresponding boson model (2.27). We note, however, that such derivation of the Jacobian for the decoupling transformation violates the $SU(2)$ symmetry, and we derive the correct expression (2.41) for the effective Lagrangian. If we eliminate $\bar{\psi}' i A'_i \tau_i \psi'$ in (2.41) by a further decoupling transformation, we will obtain the same Lagrangian as in Ref.1), although it has a much complicated form and seems to contain ghost fields. Optimistically these fields may be absorbed to the other fields and may finally disappear from the effective Lagrangian as in the ordinary Thirring model, which is open for future investigation.

References

- [1] J. Sakamoto and K. Miyake, Prog. Theor. Phys. **105** (2001), 1007.
- [2] S. Coleman, Phys. Rev. **D11** (1975), 2088.
- [3] See for example E. Abdalla, M. C. B. Abdalla and K. D. Rothe, *Non-perturbative methods in two-dimensional quantum field theory* (World Scientific, Singapore, 1991), and M. Stone, *Bosonization* (World Scientific, Singapore, 1994). References are therein.
- [4] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena* (Oxford University Press Inc., New York, 1996).
- [5] K. Fujikawa, Phys. Rev. **D21** (1980), 2848.
- [6] J. Sakamoto and B. R. Poudel, Prog. Theor. Phys. **104** (2000), 237.